

BRM MR 2635

BRL

File Cy
ADA026839

MEMORANDUM REPORT NO. 2635

A QUICK LOOK AT SPALL USING CLUSTERING
TECHNIQUES

Morton A. Hirschberg

June 1976

Approved for public release; distribution unlimited.

USA BALLISTIC RESEARCH LABORATORIES
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.
Do not return it to the originator.

Secondary distribution of this report by originating
or sponsoring activity is prohibited.

Additional copies of this report may be obtained
from the National Technical Information Service,
U.S. Department of Commerce, Springfield, Virginia
22151.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER BRL MEMORANDUM REPORT NO. 2635	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A QUICK LOOK AT SPALL USING CLUSTERING TECHNIQUES		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Morton A. Hirschberg		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS USA Ballistic Research Laboratories Aberdeen Proving Ground, MD 21005		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Materiel Development & Readiness Command 5001 Eisenhower Avenue Alexandria, VA 22333		12. REPORT DATE JUNE 1976
		13. NUMBER OF PAGES 13
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Clustering k-means ISODATA Centroid Clustering		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This report delineates the analysis of spall produced by one round of a 3.2 inch precision shaped charge using ISODATA, a centroid method clustering algorithm. Clustering was performed on the three spacial dimensions, x, y, z, by which the spall had been collected. The mass of each particle which was a priori categorized as low, medium-low, medium high, and high was then associated with its particle (by category) according to the particle's cluster membership. In general, low mass particles clustered nearer to the center of dispersion than		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. the high mass particles, a result which had been observed in earlier two dimensional regression analyses.

TABLE OF CONTENTS

	Page
I. INTRODUCTION	5
II. ANALYSIS OF SPALL DATA	6
III. DISCUSSION	9
IV. ACKNOWLEDGEMENT.	11
DISTRIBUTION LIST.	13

I. INTRODUCTION

A good portion of the data collected at the Ballistic Research Laboratories is multidimensional, consisting of spacial coordinates, velocities, accelerations, temperatures, masses, times, chemical compositions, etc.

In attempting to derive information from such data, many analytical and statistical techniques have been used. This paper delineates the use of an analytical procedure, clustering, which has had wide application in the social and biological sciences. Clustering is simply the grouping of similar objects. It is almost synonymous with classification.

As a practical example of the applicability of the technique, spall data was chosen for analysis. The spall data consists of 228 fragments of round 1198.¹ The dimensions of the problem were restricted to x,y,z coordinates (in inches), mass (in grams), and later spray angle (in degrees).

The method chosen for clustering was ISODATA,² a k-means^{2,3,4} centroid type algorithm.

Briefly, the ISODATA algorithm works as follows: a number of objects are selected as a trial subset of centroids. The entire data set is then partitioned by placing each object in the subset to which it is closest distance-wise. The average of each enlarged subset is used as the new centroid. Trial subsets are modified only after all objects have been assigned. In general, this algorithm tends to minimize the sum of the squared distances of each object from its subset mean. The user has some control over the number of clusters, as well as subset merging and splitting thresholds. In addition, the user has the option of letting the algorithm converge to a final solution, or stopping the process after a given number of iterations.

¹Rodin, B.H., Documentation of Spall Produced by a 3.2 Inch Precision Shaped Charge, Ballistic Research Laboratories, Aberdeen Proving Ground, MD, BRL Memorandum Report No. 2407, August 1974. AD #923573L.

²Rice, C.E. & Lorr, M., An Empirical Comparison of Typological Analysis Methods, ONR Contract N00014-67-A-0214, TASK0005.

³Hartigan, J. A., Clustering Algorithms, New York, Wiley 1975.

⁴Anderberg, M. R., Cluster Analysis for Applications, New York, Academic Press, 1973.

II. ANALYSIS OF SPALL DATA

The conditions under which the charge was fired, the method of collection, the masses, and the x-y coordinates are all documented in Reference 1 and will not be repeated here. The z coordinate was taken to be halfway into the sheet in which it was found (sheets were 0.5 inches thick). There was also a 24 inch offset, so the z coordinates are given by

$$(24 + 0.5(S-1) + 0.25) \quad (1)$$

where S is the sheet number. Spray angle was calculated using (2) below.

$$\text{TAN}^{-1} \frac{\sqrt{x^2+y^2}}{z} \quad (2)$$

Particle masses were divided into four groups as shown in Table I.

TABLE I
DISTRIBUTION OF PARTICLES BY MASS

<u>Mass Grams</u>	<u>Number of Particles</u>	<u>Percent of Particles</u>	<u>Group</u>
0-.002	46	20	(L) Low Mass
.002-.006	63	28	(ML) Medium-Low Mass
.006-.04	62	27	(MH) Medium-High Mass
.04 and heavier	57	25	(H) High Mass

ISODATA was run clustering the true x, y, z coordinate data of the particles. Twelve clusters were produced. The centroid data for the twelve clusters is shown in Table II.

The masses of the particles were then associated by category to the cluster to which they belonged. Table III summarizes those results.

TABLE II
CLUSTER CENTROID DATA

CLUSTER	x	y	z
1	-1.07	6.01	25.40
2	-13.91	2.45	25.36
3	-6.60	9.75	25.48
4	-2.28	13.55	25.66
5	11.40	9.12	25.55
6	5.86	5.21	25.46
7	-8.94	3.83	25.32
8	13.96	2.53	25.50
9	4.83	12.15	25.35
10	-12.81	9.19	25.75
11	-10.30	15.43	25.56
12	2.70	20.00	25.58

TABLE III
DISTRIBUTION OF PARTICLES BY MASS

CLUSTER	L(ow)	M(ed)L(ow)	MASS		Total Particles
			M(ed)H(igh)	H(igh)	
1	7	7	1	2	17
2	1	2	6	4	13
3	4	6	11	5	26
4	5	4	7	7	23
5	6	4	8	7	25
6	13	4	0	4	21
7	3	13	4	2	22
8	2	6	5	3	16
9	5	15	14	5	39
10	0	2	6	7	15
11	0	0	0	8	8
12	<u>0</u>	<u>0</u>	<u>0</u>	<u>3</u>	<u>3</u>
	46	63	62	57	228

Monte-Carlo simulations were run in order to establish the frequency with which the distribution of particles shown in Table III occurs. It was found that the distribution for clusters 1, 3, 6, 7, 9, 10, 11, 12 occurs below the chance level. That is, in general it was found that low mass particles clustered and high mass particles clustered.

Hafer⁵ showed that the mass gets larger as the spray angle increases. Mean spray angle for each cluster is shown in Table IV. In addition Table IV contains the percentage of high mass (medium-high and high) particles in each cluster.

An intuitive grouping of the clusters shows a perfect qualitative relationship between spray angle and particle mass. These results are shown in Table V.

III. DISCUSSION

A one point sample of real spall data was analyzed using ISODATA, a centroid clustering technique. Particles were found to form natural spacial clusters. When an association of particle mass with cluster membership was made, the clusters were also found to be grouped by mass. That is, in general low mass particles were found to form clusters and high mass particles were found to form clusters.

When spray angle was associated with the clusters, a perfect qualitative relationship between spray angle and particle mass was found. This result supports Hafer's work.

The work described in this report took one week to do, including writing the report. There were 3 minutes of UNIVAC 1108 computer time used in running program ISODATA and doing the Monte-Carlo runs.

It is believed that clustering techniques will be a valuable tool in the analysis of multiple burst, multiple barrel type weapons, as well as other applications (e.g. fuel fires).

⁵Hafer, A.S., An Empirically Based Scheme for Prediction of Some Characteristics of Behind-Armor Spall., Ballistic Research Laboratories, Aberdeen Proving Ground, Maryland, (Unpublished paper).

TABLE IV
MEAN SPRAY ANGLE AND PERCENTAGE OF HIGH
MASS PARTICLES BY CLUSTER

CLUSTER	MEAN SPRAY ANGLE (DEGREES)	PERCENTAGE HIGH MASS PARTICLES
1	15.76	18
2	29.09	77
3	24.88	62
4	28.23	61
5	29.74	60
6	17.71	19
7	21.80	27
8	29.15	50
9	25.80	49
10	31.73	87
11	36.20	100
12	38.76	100

TABLE V
CLUSTER GROUPING BY SPRAY ANGLE

CLUSTER GROUP	MEAN SPRAY ANGLE (DEGREES)	PERCENTAGE HIGH MASS PARTICLES
1,6	16.84	18
3,7,9	24.51	47
2,4,5,8	29.06	61
10,11,12	33.92	92

An interesting adjunct to cluster analysis would be a dimensional analysis^{6,7,8} performed first, whenever possible, to reduce the number of variables under consideration (or finding relevant variables for further analysis). Clustering (or factor analysis) applied to the reduced number of variables should provide a nice model. In addition, regression could now be performed on the resulting clusters to provide a comprehensive analysis.

Clustering therefore is useful for providing a model when none exists, quick response when little is known about the data, and pinpointing anomalies (refining a model).

It is hoped clustering techniques will be widely used in future analysis where great amounts of data exist.

IV. ACKNOWLEDGEMENT

The author would like to thank Mr. Paul Broome of the Biophysics Laboratory, Edgewood Arsenal Area, for his help in running the ISODATA program and his valuable insights into the interpretation of the results.

⁶Langhaar, H.L., Dimensional Analysis and Theory of Models, New York, Wiley, 1951.

⁷Hirschberg, M.A., A Computer Solution of the Buckingham Pi Theorem Using SYMBOLANG, A Symbolic Manipulation Language, Ballistic Research Laboratories Report No. 1824, Aberdeen Proving Ground, Maryland, August 1975. AD #A016901.

⁸Hirschberg, M.A., The Evaluation, Manipulation, and Identification of Nondimensional Numbers, Ballistic Research Laboratories, Aberdeen Proving Ground, Maryland (in preparation).

DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
12	Commander Defense Documentation Center ATTN: DDC-TCA Cameron Station Alexandria, VA 22314	2	Commander US Army Mobility Equipment Research & Development Command ATTN: Tech Docu Cen, Bldg 315 DRSME-RZT Fort Belvoir, VA 22060
1	Commander US Army Materiel Development and Readiness Command ATTN: DRCDMA-ST 5001 Eisenhower Avenue Alexandria, VA 22333	1	Commander US Army Armament Command Rock Island, IL 61202
1	Commander US Army Aviation Systems Command ATTN: DRSAV-E 12th and Spruce Streets St. Louis, MO 63166	1	Commander US Army Harry Diamond Laboratories ATTN: DRXDO-TI 2800 Powder Mill Road Adelphi, MD 20783
1	Director US Army Air Mobility Research and Development Laboratory Ames Research Center Moffett Field, CA 94035	1	Director US Army TRADOC Systems Analysis Activity ATTN: ATAA-SA White Sands Missile Range NM 88002
1	Commander US Army Electronics Command ATTN: DRSEL-RD Fort Monmouth, NJ 07703		<u>Aberdeen Proving Ground</u> Marine Corps Ln Ofc Dir, USAMSAA Cmdr, USAEA ATTN: Dr. W. Sacco Mr. P. Broome
1	Commander US Army Missile Command ATTN: DRSMI-R Redstone Arsenal, AL 35809		
1	Commander US Army Tank Automotive Development Command ATTN: DRDTA-RWL Warren, MI 48090		